

NDC 57896-180-16

**GERI CARE**

SEE NEW WARNINGS

**LIQUID  
PAIN RELIEF  
ACETAMINOPHEN****160 mg / 5 mL****ASPIRIN FREE · ALCOHOL FREE**

- **PAIN RELIEVER**
- **FEVER REDUCER**

**CHERRY FLAVOR****Compare to the active ingredient of  
Tylenol® Liquid.\*****THIS BULK PACKAGE IS NOT FOR HOUSEHOLD USE.**

Dispense contents with a child resistant closure in a tight, light-resistant container as defined in the usp.

\*This product is not manufactured or distributed by the owner of the registered trademark of Tylenol®.

**16 FL OZ (473 mL)****Drug Facts****Active ingredient (per teaspoonful = 5 mL)**  
Acetaminophen 160 mg ..... **Pain Reliever / Fever Reducer****Purpose****Uses** Temporarily relieves minor aches and pains due to: ■ Headache ■ Muscular aches ■ Backache ■ Minor pain of arthritis ■ The common cold ■ Toothache ■ Premenstrual and menstrual cramps. Temporarily reduces fever.**Warnings** **Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 6 doses in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks everyday while using this product.**Do not use:** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are allergic to acetaminophen or any of the inactive ingredients in this product.**Ask a doctor before use if the user:** ■ has a liver disease ■ is on a sodium-restricted diet**Ask a doctor or pharmacist before use if the user is:** ■ taking the blood thinning drug warfarin.**When using this product: Do not exceed recommended dose.****Stop use and ask a doctor if:** ■ Pain gets worse or lasts more than 5 days ■ Fever gets worse or lasts more than 3 days ■ New symptoms occur ■ Redness or swelling is present. These could be signs of a serious condition.**If pregnant or breast-feeding,** ask a health professional before use.**Keep out of the reach of children. Overdose warnings:** taking more than the recommended dose (overdose) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.**Directions** ■ find dose chart below, if possible, use weight to dose; otherwise use age ■ dosage may be repeated every 4 hours, or as directed by a doctor ■ do not give more than 5 doses in 24 hours.

Weight (lbs.)	under 24	24-35	36-47	48-59	60-71	72-95	over 95
Age (years)	Under 2	2 to 3	4 to 5	6 to 8	9 to 10	11 to 12	12 to adult
dosage Teaspoonful (tsp.)	consult Physician	1 tsp. (5 mL)	1 ½ tsp. (7.5 mL)	2 tsp. (10 mL)	2 ½ tsp. (12.5 mL)	3 tsp. (15 mL)	4 tsp. (20 mL)

**Other information** ■ each teaspoon (5ml) contains: sodium 7 mg. ■ store at room temperature 15° - 30°C (59°-86°F) ■ protect from freezing ■ protect from light ■ for institutional use only**Tamper Evident:** Do not use if breakaway band on cap is broken or missing.**Inactive Ingredients:** citric acid, FD & C red #40, flavor, glycerin, PEG 400, sodium benzoate, sodium citrate, sodium saccharin, sorbitol, water

DIST. BY:

Gericare Pharmaceuticals  
1650 63rd Street  
Brooklyn, NY 11204

REV. 0510R 3 57896-180-16 4